



1/20

Figure 1




2/20

Figure 2




3/20

Figure 3




4/20

Figure 4




5/20

Figure 5



6/20


Figure 6




7/20

Figure 7

A





B




8/20

Figure 8




9/20

C



D



10/20






Figure 9

8 Ångström neighbourhood

The "score" is the result of : $f(\text{subset})/f(\text{protein})$

Therefore: "scores" > 1 means that residue/group is overrepresented
"scores" < 1 means that residue/group is underrepresented

- 3,50-4,00
- 3,00-3,50
- 2,50-3,00
- 2,00-2,50
- 1,50-2,00
- 1,00-1,50
- 0,50-1,00
- 0,00-0,50



12/20


Figure 10

8 Angstrøm neighbourhood

The "score" is the result of : $f(\text{subset})/f(\text{protein})$

Therefore: "scores" > 1 means that residue/group is overrepresented

"scores" < 1 means that residue/group is underrepresented



Trp60-Cys73-Cys91

Trp118-Cys28-Cys111

Figure 11

8 Ångström neighbourhood

The triad of cutinase compared to IGG1 triads A)

The "score" is the result of : $f(\text{subset})/f(\text{protein})$

Therefore: "scores" > 1 means that residue/group is overrepresented

"scores" < 1 means that residue/group is underrepresented






Figure 12


8 Ångström neighbourhood
The triad of cutinase compared to IGG1 triads B)
The "score" is the result of : $f(\text{subset})/f(\text{protein})$

Therefore: "scores" > 1 means that residue/group is overrepresented
"scores" < 1 means that residue/group is underrepresented





15/20

Figure 13




16/20

Figure 14

A**B**

17/20


Figure 15




18/20

Figure 16

A




B



19/20

C



20/20

Figure 17 A




Figure 17B

